

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

INTERNATIONAL PATENT CLASSIFICATION⁶ : C07D 295/096, 311/68, 213/643, 405/12, A61K 31/495, 31/44, 31/35		A1	(11) International Publication Number: WO 98/00412
			(43) International Publication Date: 8 January 1998 (08.01.98)
(21) International Application Number: PCT/US97/10696			(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 26 June 1997 (26.06.97)			
(30) Priority Data: 08/674,391 1 July 1996 (01.07.96) US			
(71) Applicant: SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033 (US).			
(72) Inventors: KOZLOWSKI, Joseph, A.; P.O. Box 7391, Princeton, NJ 08543 (US). LOWE, Derek, B.; 237 North 9th Street, Kenilworth, NJ 07033 (US). CHANG, Wei, K.; 63 West Cedar Street, Livingston, NJ 07039 (US). DUGAR, Sundee; 749 Wingate Drive, Bridgewater, NJ 08807 (US).			
(74) Agents: MAGATTI, Anita, W. et al.; Schering-Plough Corporation, Patent Dept. K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).			Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

(54) Title: **MUSCARINIC ANTAGONISTS**



(57) Abstract

Di-N-substituted piperazine or 1,4-di-substituted piperidine compounds in accordance with formula (I) (including all isomers, salts, esters, and solvates), wherein Q, n, R, R¹, R², R³, R⁴, R²¹, R²⁷, R²⁸, X, and Z are as defined herein are muscarinic antagonists useful for treating cognitive disorders such as Alzheimer's disease. Pharmaceutical compositions and methods of preparation are also disclosed. Also disclosed are synergistic combinations with acetylcholinesterase inhibitors of compounds capable of enhancing acetylcholine release and having the above formula.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

MUSCARINIC ANTAGONISTS

FIELD OF THE INVENTION


The present invention relates to di-N-substituted piperazines and 1,4-di-substituted piperidines useful in the treatment of cognitive disorders, pharmaceutical compositions containing the compounds, methods of treatment using the compounds, and to the use of said compounds in combination with acetylcholinesterase inhibitors.

BACKGROUND OF THE INVENTION

Alzheimer's disease and other cognitive disorders have received much attention lately, yet treatments for these diseases have not been very successful. According to Melchiorre *et al.* (*J. Med. Chem.* (1993), 36, 3734-3737), compounds that selectively antagonize m2 muscarinic receptors, especially in relation to m1 muscarinic receptors, should possess activity against cognitive disorders. Baumgold *et al.* (*Eur. J. Pharmacol.*, 251, (1994) 315-317) disclose 3- α -chloroimperialine as a highly selective m2 muscarinic antagonist.

The present invention is predicated on the discovery of a class of di-N-substituted piperazines and 1,4-di-substituted piperidines. Logemann *et al.* (*Brit. J. Pharmacol.* (1961), 17, 286-296) describe certain di-N-substituted piperazines, but these are different from the inventive compounds of the present invention. Furthermore, the compounds of Logemann *et al.* are not disclosed to have activity against cognitive disorders.

International Patent Publication Number WO93/08799 published May 13 1993 (Smith-Kline Beecham) discloses *inter alia* indane derivatives that are endothelin receptor antagonists and are (in part) of the following formula:



wherein:

R_1 is $-X(CH_2)_nAr$ or $-X(CH_2)_nR_8$;

R_2 is H or Ar;

P_1 is $-X(CH_2)_nR_8$;

P_2 is $-X(CH_2)_nR_8$ or $-XR_9Y$;

R_8 is H, alkyl, alkenyl, alkynyl, CO_2H , CO_2alkyl , or CO_2Ar ;

- 2 -

R₉ is alkyl, alkenyl or phenyl;

R₁₀ is H, alkyl (which may be substituted with CO₂H, CO₂alkyl or CO₂(CH₂)_nAr), alkenyl, phenyl, OH, alkoxy, S(O)_qalkyl, S(O)_qalkenyl, S(O)_qaryl, NH₂, NHalkyl, N(alkyl)₂, F, Cl, Br, I, CF₃, NHCHO, NHCOalkyl, -X(CH₂)_nR₈ or
 5 -XR₉Y;

X is (CH₂)_n, O, NH, Nalkyl, or S(O)_q;

Y is CH₃ or X(CH₂)_nAr;

Ar is a variety of substituted or unsubstituted heterocyclic and aromatic hydrocarbon groups, including piperidiny and piperazinyl, which may carry
 10 substituents;

Z₁ and Z₂ are independently H, alkyl, alkenyl, alkynyl, OH, alkoxy, S(O)_qalkyl, NH₂, NHalkyl, N(alkyl)₂, F, Cl, Br, I, CF₃, NHCHO, NHCOalkyl, -X(CH₂)_nR₈, phenyl, benzyl or cycloalkyl;

Z₃ is Z₁ or -XR₉Y;


15 n is 0 or an integer from 1 to 6, and q is 0, 1 or 2;

and the groups designated as 'alkyl', 'alkenyl', 'alkynyl' or 'phenyl' can all be substituted.

(The definitions of radicals as given in that patent do not in general pertain to the present invention, even though similar symbols may be used.)

20 SUMMARY OF THE INVENTION

The present invention relates to compounds according to the structural formula I,



I

including all stereoisomers and pharmaceutically acceptable salts, esters,
 25 and solvates thereof,
 wherein:


Z is N, CH or C-alkyl;

X is -O-, -S-, -SO-, -SO₂-, -CO-, -CH₂-, -CONR²⁰-, -NR²⁰-SO₂-,
 -NR²⁰CO-, or -SO₂-NR²⁰-;

30 Q is -O-, -S-, -SO-, -SO₂-, or -CH₂-;


- 3 -

R is



R¹ and R²¹ are independently selected from the group consisting of hydrogen, alkyl, alkenyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl, cycloalkenylalkyl, phenylalkyl, and hydroxyalkyl;

R² is cycloalkyl, cycloalkyl substituted with 1 to 3 independently selected R³ groups, cycloalkenyl, cycloalkylalkyl,



(wherein R^γ is H, alkyl, alkenyl, SO₂R^z or COR^z wherein R^z is alkyl, alkenyl, aryl, heteroaryl, or cycloalkyl), with the proviso that R² is R³-substituted-1-piperidiny1 only when Z is CH or C-alkyl; or, when Z is CH, R² may also be alkoxycarbonyl, -N(R⁹)(hydroxyalkyl) wherein R⁹ is H, hydroxyalkyl, or alkyl, or -N(R⁹)₂ wherein the two R⁹ groups may be joined to form an alkylene group;

R³, R⁴, R⁵, R⁶, R²², R²⁴, and R²⁵ are independently selected from the group consisting of H, alkyl, halo, alkoxy, benzyloxy, benzyloxy substituted by nitro or aminoalkyl, polyhaloalkyl, nitro, sulfonyl, hydroxy, amino, alkylamino, formyl, alkylthio, acyloxy, alkylsulfonyl, arylsulfonyl, acyl, alkoxycarbonyl, alkylsulfinyl, -OCONH₂, -OCONH-alkyl, -OCON(alkyl)₂, -NHCOO-alkyl, -NHCO-alkyl, phenyl, hydroxyalkyl, and 1-morpholinyl;

R⁸ is hydrogen, lower alkyl or cyclopropyl;

R²⁰ is H, phenyl or alkyl;

R²⁷ and R²⁸ are independently selected from the group consisting of H, alkyl, hydroxyalkyl, alkoxyalkyl, arylalkyl, mercaptoalkyl, alkylthioalkyl, and carboxyalkyl, and additionally R²⁷ and R²⁸ may be joined to form an alkylene group; and

n is 0 or an integer from 1 to 3.

Another aspect of the invention is a pharmaceutical composition which comprises a compound having structural formula I as defined above, including

- 4 -

stereoisomers, pharmaceutically acceptable salts, esters, and solvates thereof, in combination with a pharmaceutically acceptable carrier.

Another aspect of the invention is the use of a compound of formula I as defined above, including stereoisomers, pharmaceutically acceptable salts, esters, and solvates thereof, for the preparation of a pharmaceutical composition useful in the treatment of cognitive disorders and neurodegenerative diseases such as Alzheimer's disease.

Yet another aspect of the invention comprises a method for making a pharmaceutical composition comprising mixing a compound of formula I as defined above, including stereoisomers, pharmaceutically acceptable salts, esters, and solvates thereof, with a pharmaceutically acceptable carrier.

Another aspect of this invention is a method for treating a cognitive or neurodegenerative disease comprising administering to a patient suffering from said disease an effective amount of a compound of formula I as defined above, including stereoisomers, pharmaceutically acceptable salts, esters, and solvates thereof.

Another aspect of this invention is a method for treating cognitive and neurodegenerative diseases, such as Alzheimer's disease, with a compound of formula I as defined above, including stereoisomers, pharmaceutically acceptable salts, esters, and solvates thereof, in combination with an acetylcholinesterase inhibitor.

Another aspect of this invention is a method for treating a cognitive or neurodegenerative disease comprising administering to a patient suffering from said disease an effective amount of a combination of a compound of formula I as defined above, including stereoisomers, pharmaceutically acceptable salts, esters, and solvates thereof, said compound being capable of enhancing acetylcholine release (and being preferably an m2 or m4 selective muscarinic antagonist), together with an acetylcholinesterase inhibitor.

Another aspect of this invention is a kit comprising in separate containers in a single package pharmaceutical compounds for use in combination to treat cognitive disorders, wherein

one container contains a compound of formula I as defined above, including stereoisomers, pharmaceutically acceptable salts, esters, and solvates thereof, said compound being capable of enhancing acetylcholine release (and preferably being an m2 or m4 selective muscarinic antagonist) in a pharmaceutically acceptable carrier, and

- 5 -

a second container contains an acetylcholinesterase inhibitor in a pharmaceutically acceptable carrier,
the combined quantities being an effective amount.


DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

5 In a particularly preferred group of compounds of the formula I, Z is N. In another preferred group, n is 1 or 2 or especially 0.

In another preferred group of compounds, X is SO or especially O or SO₂.

In yet another preferred group of compounds, R⁸ is H or methyl.

In another preferred group of compounds, R is

10  (wherein R⁵ and R⁶ are H, CH₃, nitro, NH₂, acetylamino, or methoxy), especially 4-methoxyphenyl; X is preferably O, SO or SO₂, R³ and R⁴ are H, and R¹ is H, cycloalkyl, cycloalkylalkyl or alkyl, and R²¹ is H.

15 In another preferred group of compounds, R³ and R⁴ are H, and R¹ is H, cycloalkyl or alkyl and R²¹ is H. R¹ is preferably H, CH₃ or cyclohexylmethyl, and R² is cyclohexyl.

In another preferred group of compounds, R³ and R⁴ are H, X is O, SO or SO₂, and R¹ is H, cycloalkyl or alkyl and R²¹ is H. R¹ is preferably H or CH₃.

20 In another preferred group of compounds, at least one of R²⁷ and R²⁸ is alkyl. In particular, at least one of R²⁷ and R²⁸ is alkyl and the other is H or alkyl; more preferably, one of R²⁷ and R²⁸ is methyl and the other is hydrogen.

In another preferred group of compounds, R is 4-methoxyphenyl.

In a particularly preferred group of compounds of the formula I:

25 R is a phenyl group, which may be substituted with nitro or especially with methoxy, where each of these groups is preferably in the 4-position, or in particular a 2-pyrimidinyl group;

X is S, SO, or especially SO₂ or O;

Q is O or especially CH₂;

n is 1 or 2 or especially 0;

30 R¹ is H and R²¹ is cyclohexylmethyl;

R²⁷ and R²⁸ are CH₃ or especially H;

Z is N; and

R² is cyclohexyl or 1-piperidinyl.

- 6 -

Except where stated otherwise the following definitions apply throughout the present specification and claims. These definitions apply whether a term is used by itself or in combination with other terms. For example, the definition of "alkyl" applies not only to "alkyl" but also to the "alkyl" portions of "alkoxy",

5 "polyhaloalkyl", etc.

Alkyl represents a straight or branched saturated hydrocarbon chain having 1 to 20 carbon atoms, preferably 1 to 8 carbon atoms. A 'lower alkyl' group has 1 to 6 carbon atoms, preferably 1 to 4 carbon atoms.

10 Alkenyl represents a straight or branched hydrocarbon chain of from 3 to 15 carbon atoms, more preferably 3 to 12 carbon atoms, having at least one carbon-to-carbon double bond but the free valency at least one carbon atom removed from the double bond.

Cycloalkyl represents a saturated carbocyclic ring having 3 to 12 carbon atoms.

15 Cycloalkenyl represents a carbocyclic ring having from 5 to 8 carbon atoms and at least one carbon-to-carbon double bond in the ring.

Acyl represents a radical of a carboxylic acid and thus includes groups of the formula Alkyl-CO-, Aryl-CO-, Aralkyl-CO-, Cycloalkyl-CO-, wherein the various hydrocarbon radicals are as defined in this section.

20 Halo represents fluoro, chloro, bromo or iodo.

Aryl represents phenyl or naphthyl, each of which may be substituted with one to three groups R^c selected from halo, alkyl, hydroxy, alkoxy, phenoxy, amino, alkylamino and dialkylamino groups. Preferred aryl groups are phenyl, substituted phenyl, 1-naphthyl, 2-naphthyl and indanyl groups.

25 Heteroaryl represents a cyclic group having at least one O, S and/or N interrupting a carbocyclic ring structure and having a sufficient number of pi electrons to provide aromatic character. The aromatic heterocyclic group preferably has from 2 to 14, especially from 3 to 9 carbon atoms, e.g., 2-, 3- or 4-pyridyl, 2- or 3-furyl, 2- or 3-thienyl, 2-, 4- or 5-thiazolyl, 2-, 4- or 5-imidazolyl, 30 4-, 5- or especially 2-pyrimidinyl, 2-pyrazinyl, 3- or 4-pyridazinyl, 3-, 5- or 6-[1,2,4-triazinyl], 3- or 5-[1,2,4-thiadiazolyl], 2-, 3-, 4-, 5-, 6- or 7-benzofuranyl, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 3-, 4- or 5-pyrazolyl, or 2-, 4- or 5-oxazolyl, etc. Preferred heteroaryl groups include 2-, 3- or 4-pyridyl, 2- or 3-furyl, 2- or 3-thienyl, 2-, 4- or 5-imidazolyl, and 7-indolyl.

35 Polyhalo indicates substitution of at least 2 halo atoms in the group modified by the term "polyhalo".

Sulfonyl represents a group of the formula -SO₂-.


- 7 -

Sulfinyl represents a group of the formula $-\text{SO}-$.

Alkylene represents a straight or branched saturated hydrocarbon chain having 1 to 20 carbon atoms, preferably 1 to 8 carbon atoms, and two free valencies, which for the purpose of this invention are not on the same carbon atom when the alkylene group has 2 to 20 carbon atoms. Preferred alkylene groups are methylene or polymethylene groups of the formula $-(\text{CH}_2)_{(2-20)}-$.

Each radical or group that appears more than once in a structural formula, for example R^9 when R^2 is $-\text{N}(\text{R}^9)_2$, may be independently selected from the whole definition for that radical or group.

Compounds of this invention may exist in at least two stereoisomeric configurations based on the asymmetrically substituted carbon atom to which



the group R^2 is attached. Further stereoisomerism may be present, for example when R^1 and R^{21} are not identical, or when X is SO , or when at least one of R^{27} and R^{28} is not hydrogen. All possible stereoisomers of formula I are within the scope of the invention.

Compounds of formula I can exist in unsolvated as well as solvated forms, including hydrated forms. In general, the forms that are solvated with pharmaceutically acceptable solvents such as water, ethanol and the like, are equivalent to the unsolvated forms for the purposes of this invention.


A compound of formula I may form pharmaceutically acceptable salts with organic and inorganic acids. Examples of suitable acids for salt formation are hydrochloric, sulfuric, phosphoric, acetic, citric, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic and other mineral and carboxylic acids well known to those skilled in the art. A salt is prepared by contacting a free base form with a sufficient amount of the desired acid to produce a salt in the conventional manner. The free base form may be regenerated by treating the salt with a suitable dilute aqueous base such as dilute aqueous sodium hydroxide, potassium carbonate, ammonia or sodium bicarbonate. The free bases differ from the corresponding salts somewhat in certain physical properties, such as solubility in polar solvents, but the salts are otherwise equivalent to the corresponding free bases for purposes of the invention.

Compounds of formula I and their salts can be prepared by standard methods. The methods of the following Schemes are preferred; the starting materials either are known or can be prepared by standard methods, and the

- 8 -

radicals in the formulae (unless otherwise stated) have the meanings given for formula I, with the proviso that R^2 can also be a nitrogen-protecting group that is replaced (e.g., at the end of the Scheme) with a group R^2 according to formula I:

Scheme A:



5

In Scheme A, Q can be $-O-$, $-S-$, or $-CH_2-$. In Step (a), L^1 and L^2 are groups that can be eliminated during the reaction. L^2 is preferably a leaving group such as a halogen atom, especially bromine or fluorine, and L^1 is preferably an alkali metal, e.g., sodium or potassium. The reaction is carried out in an inert organic solvent, which is preferably anhydrous, e.g., DMSO, DMF, or a polar ether such as dimethoxyethane.

10

When X is O or S, the reactant contributing the group X can be prepared by reaction of the parent compound containing the group XH and an alkali metal hydride *in situ* in the inert organic solvent.

15


Step (b) comprises the reduction of the carbonyl group to hydroxymethylene with a reducing agent such as an aluminum hydride, e.g., lithium

- 9 -

aluminum hydride, or a borohydride, e.g., sodium borohydride, in an inert organic solvent. When an aluminum hydride is used, the solvent should be anhydrous and is preferably an ether solvent such as THF; when a borohydride is used, the solvent can be a lower alkanol, especially methanol or ethanol, or THF, DMF, or DMSO, all of which may be anhydrous or aqueous. Step (c) comprises the conversion of the hydroxy group into a leaving group L³, e.g., a sulfonate ester group such as a group SO₃-(lower alkyl) or with a halogen atom, especially bromine or iodine. This conversion can be carried out with a sulfonyl chloride, e.g., methane-, ethane- or 4-toluene-sulfonylchloride, and an organic base, e.g., a tertiary amine such a pyridine or triethylamine; or with a halogenating agent, e.g., SOCl₂ or PBr₃, if desired with displacement of the resulting bromide by iodide when an iodide is required.

Step (d) comprises the amination of the reactive group L³, e.g. Cl, with a piperidine or piperazine to form the desired product of the formula I. This reaction can be carried out neat with the piperidine or piperazine or in an organic solvent, and an acid-binding agent can if desired be used. An excess of the piperidine or piperazine may serve both as acid-binding agent and as organic solvent. When a piperazine is used, its second nitrogen atom can bind the acid liberated in the reaction and no extra acid-binding agent may be necessary.

Step (e) can be carried out by condensation of the reactants in the presence of a basic catalyst, e.g, an amine such as piperidine. Under these circumstances an excess of the catalyst can serve as solvent. Alternatively, the reaction can be effected in the presence of a strongly basic catalyst such as LDA (lithium diisopropylamide) or lithium HMDS (lithium hexamethyldisilazane), which preferably is added at a low temperature, e.g., -78°C to 0°C, in an inert anhydrous organic solvent such as THF or diethyl ether. The reaction with the aldehyde may then be effected at a low temperature, at room temperature or at moderately elevated temperature, e.g., at -78 to +60°C, preferably about 0 to +30°C, and under an inert atmosphere, e.g., nitrogen. When the aldehyde R²¹A-CHO is used to introduce the group R²¹, then R²¹A has one carbon atom fewer than R²¹ such that R²¹A-CH₂ is R²¹. The intermediate carbinol of the formula



- 10 -

can then be dehydrated (for example, direct with acid or, after conversion of the hydroxy group into a leaving group, e.g., a halide or sulfonate, with base), and the resulting compound with a double bond can be hydrogenated, e.g. with hydrogen and a catalyst such as Pd/C, to the compound of the formula A6.

- 5 When the hydrogenation also reduces the carbonyl group, then the reduction of the following Step (f) can be obviated.


Step (f) can then be carried out by the methods of Steps (b) (reduction) and (c) (conversion of the hydroxy group into a leaving group L³) above, and Step (g) by the method of Step (d) above.

- 10 If it is desired to introduce a group R¹ in addition to R²¹, then the process of Step (e) can be repeated before Steps (f) and (g) are carried out.

- An alternative method for preparing compounds of the formulae A5 and A7 in Scheme A comprises the condensation of a compound of the formula A2 or A6 with a compound of the formula A4, followed by the reduction of the resulting condensation product, e.g., imine, preferably in the same step. The condensation can be effected in the presence of a compound or compounds serving as a mild Lewis acid (or a protic acid) and a dehydrating agent. The mild Lewis acid is conveniently a titanium tetra(lower alkoxide), especially Ti(O-2-Pr)₄, which is commercially available and gives good results. The resulting condensation product (e.g., imine) can then be reduced with a mild reducing agent, e.g., a sodium borohydride, but preferably one that is not reactive towards the Lewis acid or protic acid, especially sodium cyanoborohydride. Alternatively, the reaction can be effected with sodium triacetoxyborohydride as both Lewis acid and reducing agent, in the presence of acetic acid.
- 15
20
25

The condensation and reduction is carried out in the presence of an inert organic solvent, preferably a chlorinated hydrocarbon, e.g., 1,2-dichloroethane or especially methylene chloride.

- 11 -

Scheme B


In Scheme B, Q is -O- or -S-. In Step (a) the compound of the formula B1 (wherein R^A is R or preferably H) is reacted with a reactive derivative of an acid of the formula L⁴·(CH₂)_n·CH₂·CO₂H, or such an acid with a substituent R¹ and/or R²¹ on the 2-carbon atom, wherein L⁴ is a reactive group such as a chlorine or bromine atom. The reactive derivative is preferably the acid chloride, but may be the bromide or anhydride. The reaction is effected under Friedel-Crafts conditions with a catalyst and inert organic solvent; the catalyst is for example anhydrous aluminum chloride or boron trifluoride, and the solvent is preferably nitrobenzene. The reaction is typically started at low temperature, e.g., at 0–20°C, and continued at moderately elevated temperature, e.g., 30–100°C.

A group R²¹ (where R²¹ is other than hydrogen) may be introduced in Step (b) as described under Scheme A above. When the group R^A is hydrogen, the group R may be introduced in Step (c) by reaction of the starting material with a compound of the formula RL⁴, wherein L⁴ is a leaving group, e.g., a sulfonate ester group but preferably a halogen, especially Cl or Br. The reaction is preferably carried out in an organic solvent in the presence of a strong

- 12 -

inorganic base, e.g., sodium or potassium hydride or hydroxide, and an inert organic solvent, e.g. anhydrous DMSO, DMF, or a polar ether such as dimethoxyethane. Steps (d) and (e) may then be carried out as described under Scheme A above for Steps (b), (c) and (d) therein.

5 Scheme C:




In Scheme C, Q is O or S. In Step (a), the group R is introduced by reaction of the starting material with a compound of the formula RL^4 , wherein L^4 is a leaving group, as described under Step (c) of Scheme B above. The remaining steps (b) through (g) can then be effected according to the processes described above for steps (b) through (g) respectively of Scheme A.

10

- 13 -


Scheme D:



- The process of Step (a) (reductive amination) can be carried out according to the alternative condensation/reduction process of Scheme A (described above as an alternative method for preparing compounds of the formulae A5 and A7). The group L^5 is preferably a chlorine atom or especially a bromine or iodine atom. The product of the formula D2 can then be reacted with an alkyl-lithium reagent, e.g., *n*-BuLi or *t*-BuLi, in an inert anhydrous organic solvent, to replace the group L^5 with Li, and the resulting organometallic compound can be reacted with the aldehyde RCHO to yield the compound of the formula D3. The benzylic hydroxy group in D3 can then be reduced, for example with a trialkylsilane, preferably Et_3SiH , and a strong acid, preferably trifluoroacetic acid. If the product of the formula D4 is to contain a group R^1 or R^{21} , this group may be introduced into the compound of the formula D1.
- The group R^2 in this process is preferably a nitrogen-protecting group, e.g., a benzyloxycarbonyl (CBZ) or especially a *t*-butyloxycarbonyl (BOC) group. Such a protecting group can be replaced with a group R^2 defined above under formula I. A protecting group such as an amidating group can be removed by hydrolysis; e.g., a protecting BOC group can be removed with trifluoroacetic acid or with HCl in EtOAc. The desired group R^2 can then be introduced (in one

- 14 -

or more steps); for example, a 1-piperidinyl group or N-RY-1-piperidinyl group can be introduced by reductive condensation with a compound of the formula



5 followed by removal of the protecting BOC group by hydrolysis. Both the reductive condensation and the hydrolysis can be effected as described above. Any further substituent RY (wherein RY is as defined above) can then be introduced on to the newly liberated nitrogen atom. For example, an amide can be formed by condensation with the acid chloride (RY.Cl, when RY is an acyl group), if necessary in the presence of a base, or by condensation with the acid
10 itself (RY.OH, when RY is an acyl group) in the presence of 1-hydroxy-benzotriazole, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide, and N-methyl-morpholine.

After the process of Scheme A, B, C or D has been carried out, a resulting compound of the formula I wherein Q represents S can be oxidized to a
15 compound wherein Q represents SO or SO₂; the oxidizing agent is preferably a peracid such as peracetic acid, 3-chloroperbenzoic acid, or perboric acid, in the presence of a strong acid used in excess, e.g., methanesulfonic acid. If the compound wherein Q is S is being oxidized to the sulfoxide (or the sulfoxide to the sulfone), then about one equivalent of oxidizing agent should be used; if the
20 compound wherein Q is S is being oxidized to the sulfone, then about two equivalents (preferably at least two equivalents) of oxidizing agent should be used.

PHARMACOLOGICAL ACTIVITY

25 The compounds of formula I exhibit selective m₂ and/or m₄ muscarinic antagonizing activity, which has been correlated with pharmaceutical activity for treating cognitive disorders such as Alzheimer's disease and senile dementia.

The compounds of formula I display pharmacological activity in test procedures designed to indicate m₁, m₂ and m₄ muscarinic antagonist activity. The compounds are non-toxic at pharmaceutically therapeutic doses. Following
30 are descriptions of the test procedures.

MUSCARINIC BINDING ACTIVITY

The compound of interest is tested for its ability to inhibit binding to the cloned human m₁, m₂, and m₄ muscarinic receptor subtypes. The sources of receptors in these studies were membranes from stably transfected CHO cell
35 lines which were expressing each of the receptor subtypes. Following growth,

- 15 -

the cells were pelleted and subsequently homogenized using a Polytron in 50 volumes cold 10 mM Na/K phosphate buffer, pH 7.4 (Buffer B). The homogenates were centrifuged at 40,000 x g for 20 minutes at 4°C. The resulting supernatants were discarded and the pellets were resuspended in
5 Buffer B at a final concentration of 20 mg wet tissue/ml. These membranes were stored at -80°C until utilized in the binding assays described below.

Binding to the cloned human muscarinic receptors was performed using ³H-quinuclidinyl benzilate (QNB) (Watson et al., 1986). Briefly, membranes (approximately 8, 20, and 14 µg of protein assay for the m1, m2, and
10 m4-containing membranes, respectively) were incubated with ³H-QNB (final concentration of 100-200 pM) and increasing concentrations of unlabeled drug in a final volume of 2 ml at 25°C for 90 minutes. Non-specific binding was assayed in the presence of 1 µM atropine. The incubations were terminated by vacuum filtration over GF/B glass fiber filters using a Skatron filtration apparatus,
15 and the filters were washed with cold 10 mM Na/K phosphate buffer, pH 7.4. Scintillation cocktail was added to the filters, and the vials were incubated overnight. The bound radioligand was quantified in a liquid scintillation counter (50% efficiency). The resulting data were analyzed for IC₅₀ values (i.e. the concentration of compound required to inhibit binding by 50%) using the EBDA
20 computer program (McPherson, 1985). Affinity values (K_i) were then determined using the following formula (Cheng and Prusoff, 1973):

$$K_i = \frac{IC_{50}}{1 + \left[\frac{\text{concentration of radioligand}}{\text{affinity (K}_D\text{) of radioligand}} \right]}$$

Hence a lower value of K_i indicates greater binding affinity.

The above procedure is known in the art and has been the subject of
25 detailed publications.

To determine the degree of selectivity of a compound for binding the m2 receptor, the K_i value for m1 receptors was divided by the K_i value for m2 receptors. A higher ratio indicates a greater selectivity for binding the m2 muscarinic receptor.

30 MICRODIALYSIS METHODOLOGY

The following procedure is used to show that a compound functions as an m2 antagonist:

Surgery: For these studies, male Sprague-Dawley Rats (250-350 g) were anesthetized with sodium pentobarbital (54 mg/kg, ip) and placed on a
35 Kopf stereotaxic apparatus. The skull was exposed and drilled through to the

- 16 -

dura at a point 0.2 mm anterior and 3.0 mm lateral to the bregma. At these coordinates, a guide cannula was positioned at the outer edge of the dura through the drilled opening, lowered perpendicularly to a depth of 2.5 mm, and permanently secured with dental cement to bone screws. Following the surgery, rats were given ampicillin (40 mg/kg, ip) and individually housed in modified cages. A recovery period of approximately 3 to 7 days was allowed before the microdialysis procedure was undertaken.

Microdialysis: All of the equipment and instrumentation used to conduct *in vivo* microdialysis was obtained from Bioanalytical Systems, Inc. (BAS). The microdialysis procedure required the insertion through the guide cannula of a thin, needle-like perfusable probe (CMA/12.3 mm x 0.5 mm) to a depth of 3 mm in striatum beyond the end of the guide. The probe was connected beforehand with tubing to a microinjection pump (CMA-/100). Rats were collared and tethered, and, following probe insertion, were placed in a large clear plexiglass bowl with litter material and access to food and water. The probe was perfused at 2 µl/min with Ringer's buffer (NaCl 147 mM; KCl 3.0 mM; CaCl₂ 1.2 mM; MgCl₂ 1.0 mM) containing 5.5 mM glucose, 0.2 mM L-ascorbate, and 1 µM neostigmine bromide at pH 7.4. To achieve stable baseline readings, microdialysis was allowed to proceed for 90 minutes prior to the collection of fractions. Fractions (20 µl) were obtained at 10-minute intervals over a 3-hour period using a refrigerated collector (CMA/170 or 200). Following the collection of four to five baseline fractions, the drug or combination of drugs to be tested was administered to the animal. Once collection was complete, each rat was autopsied to determine how accurately the probe was placed.

Acetylcholine (ACh) analysis: The concentration of ACh in collected samples of microdialysate was determined using HPLC/electrochemical detection. Samples were auto-injected (Waters 712 Refrigerated Sample Processor) onto a polymeric analytical HPLC column (BAS, MF-6150) and eluted with 50 mM Na₂HPO₄, pH 8.5. To prevent bacterial growth, Kathon CG reagent (0.005%) (BAS) was included in the mobile phase. Eluate from the analytical column, containing separated ACh and choline, was then immediately passed through an immobilized enzyme reactor cartridge (BAS, MF-6151) coupled to the column outlet. The reactor contained both acetylcholinesterase and choline oxidase covalently bound to a polymeric backbone. The action of these enzymes on ACh and choline resulted in stoichiometric yields of hydrogen peroxide, which was electrochemically detected using a Waters 460 detector equipped with a platinum electrode at a working potential of 500 mvolts. Data

- 17 -

acquisition was carried out using an IBM Model 70 computer equipped with a microchannel IEEE board. Integration and quantification of peaks were accomplished using "Maxima" chromatography software (Waters Corporation). Total run time per sample was 11 minutes at a flow rate of 1 ml/min. Retention times for acetylcholine and choline were 6.5 and 7.8 minutes respectively. To monitor and correct for possible changes in detector sensitivity during chromatography, ACh standards were included at the beginning, middle and end of each sample queue.

Increases in ACh levels are consistent with presynaptic m2 receptor antagonism.

RESULTS OF THE TESTS

The present invention also relates to achieving similar synergistic results by administering any compound capable of enhancing ACh release, such as a muscarinic antagonist, e.g., scopolamine or QNB, in combination with an acetylcholinesterase inhibitor. Preferably the ACh release-enhancing compound is either an m2 selective muscarinic antagonist, i.e. one having a ratio of $(K_i \text{ for m1})/(K_i \text{ for m2})$ greater than 1, or an m4 selective muscarinic antagonist, i.e. one having a ratio of $(K_i \text{ for m1})/(K_i \text{ for m4})$ greater than 1. The m2 or m4 selective muscarinic antagonists for practicing this aspect of the invention include without limitation 3- α -chloroimperialine, AF-DX 116, AF-DX 384, BIBN 99 (these last three compounds being available from Boehringer-Ingelheim), tripitramine, and himbacine.

The tests reported in the following Table show that the compounds numbered 5, 9, 10, 14 and 15 in particular have useful values of the m1/m2 and m4/m2 ratios and should have valuable properties for the treatment of cognitive disorders.

30



35

- 18 -

TABLE 1: ACTIVITIES OF COMPOUNDS OF FORMULA I	Compound No.	M.P. °C	K _i (nm); m1/m2 & m4/m2			
			m1 $\frac{m1}{m2}$	m2	m4 $\frac{m4}{m2}$	
	1		356 4.5	79	106 1.3	
	2	67-68	42 1.5	27		
	3	219-221	82 0.6	130	37 0.3	
	4	124-126	223 1.0	224	>200 >0.9	
	5	179-181, dimaleate salt	160 7.5	21	35 1.6	
	6	205 (dec.)	633 6.6	96	77 0.8	
	7	191-193, HCl salt	105 4.0	26	21 0.8	
	8	191-193, HCl salt	88 3.3	27	23 0.9	
	9	204-206, dimaleate salt	265 7.7	35		
	10	183-184, HCl salt	151 13.2	11.5	17 1.5	
	11	181-182, HCl salt	314 3.6	88		
	12	150-154, dimaleate salt	173 1.9	92		
	13		67 4.0	17	33 2.0	
	14 (Isomer A)	232, HCl salt	112 18.4	6.1	16 2.6	
	15 (Isomer B)	253-254, HCl salt	1.23 11.2	0.11	0.27 2.5	

- 19 -

TABLE 2: Effect of Compounds of the Invention on Release of Acetyl Choline (ACh) from Striatum of Conscious Rats following Peritoneal Administration


Sample Collection Time (Minutes)	Compound and results		Compound and results	
				
	Compound 14, Isomer A		Compound 15, Isomer B	
	Dose = 10 mg/kg; 3 rats		Dose = 10 mg/kg; 3 rats	
	Mean % of baseline	Standard error	Mean % of baseline	Standard error
20 (1)	99.47	0.83	98.61	1.74
30 (2)	100.85	4.62	93.87	0.13
40 (3)	99.71	3.14	102.42	1.41
50 (4)	99.20	1.59	103.86	3.90
60	100.78	4.20	102.14	4.10
70	86.66	6.26	175.31	6.60
80	95.26	4.15	278.67	31.06
90	95.90	1.66	266.94	17.84
100	102.30	16.27	245.19	16.74
110	108.24	13.56	217.34	9.29
120	85.62	16.35	222.39	20.20
130	103.94	13.20	205.69	12.36
140	99.35	13.94	188.49	11.52
150	91.47	11.51	189.79	13.56
160	90.80	20.28	184.27	13.03
170	89.0	12.35	189.27	23.70
180	83.13	10.95	189.86	12.41

Notes:


- 5 (1) Control; gives first baseline.
 (2) Control; gives second baseline.
 (3) Control; gives third baseline.
 (4) Compounds were injected at 50 minutes.

- 20 -




Particularly preferred compounds of the formula I include those of the following formulae, and their acid addition salts:




and



and also the compounds of the following formulae, and their acid addition salts:



and



5

in all of these compounds, the methyl group attached to the piperazine ring (where present) is in the (R)-conformation.

Pharmaceutical compositions can be prepared from the compounds of formula I, which are capable of enhancing ACh release, by admixing them with pharmaceutically acceptable, inert carriers. Acetylcholinesterase inhibitors can be used as optional constituents of such pharmaceutical compositions to provide a better, frequently even synergistic, effect. The pharmaceutically acceptable carriers may be either solid or liquid. Preparations in solid form include powders, tablets, dispersible granules, capsules, cachets and

10

- 21 -

suppositories. A solid carrier can be one or more substances which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders or tablet-disintegrating agents; it may also be an encapsulating material.

5 Preparations in liquid form include solutions, suspensions and emulsions. As an example may be mentioned solutions in water or in water-propylene glycol for parenteral injection.

Also included are preparations in solid form which are intended for conversion, shortly before use, to liquid form preparations for either oral or
10 parenteral administration. Such liquid forms include solutions, suspensions and emulsions. The solid forms are most conveniently provided in unit dose form for this purpose and are used to provide a single liquid dosage unit.

The invention also contemplates alternative delivery systems including, but not necessarily limited to, transdermal delivery. The transdermal
15 compositions can take the form of creams, lotions and/or emulsions and can be included in transdermal patches of the matrix or reservoir type as are conventional in the art for this purpose.

Preferably, the pharmaceutical preparation is in unit dosage form. In such form, the preparation is subdivided into unit doses containing appropriate
20 quantities of the active components. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation such as packeted tablets, capsules and powders in vials or ampoules. The unit dosage form can also be a capsule, cachet or tablet itself, or it may be the appropriate number of any of these in a packaged form.

25 The quantity of active compound in a unit dose preparation may be varied or adjusted from 1 mg to 100 mg according to the particular application and the potency of the active ingredient and the intended treatment. This would correspond to a dose of about 0.001 to about 20 mg/kg, which may be divided over 1 to 3 administrations per day. The composition may, if desired, also
30 contain other therapeutic agents.


The dosages may be varied, depending on the requirement of the patient, the severity of the treated condition, and the particular compound administered. Determination of the proper dosage for a particular situation is within the skill of those in the medical art. For convenience, the total daily dosage may be divided
35 and administered in portions throughout the day or by means providing continuous delivery.

- 22 -


When a compound of formula I or a compound capable of enhancing ACh release is used in combination with an acetylcholinesterase inhibitor to treat cognitive disorders, these two active components may be co-administered simultaneously or sequentially. Alternatively, a single pharmaceutical composition comprising a compound of formula I or a compound capable of enhancing ACh release and an acetylcholinesterase inhibitor in a pharmaceutically acceptable carrier can be administered. The components of the combination can be administered individually or together in any conventional oral or parenteral dosage form, such as capsule, tablet, powder, cachet, suspension, solution, suppository, nasal spray, etc. The dosage of the acetylcholinesterase inhibitor may range from 0.001 to 100 mg/kg body weight.

The invention disclosed herein is exemplified by the following Examples, which should not be construed to limit the scope of the disclosure. Alternative mechanistic pathways and analogous structures may be apparent to those skilled in the art.

EXAMPLES

Example 1:Step A


4.2 g (29 mmoles) of 5-fluoroindanone and 5.05 g (31 mmoles) of sodium benzenesulfonate were combined in dry DMSO (20 mL). The mixture was heated to about 120°C for 48 hours, cooled to room temperature, and poured into ice water (300 mL). The precipitate that developed was filtered off and triturated with 100 mL dry ether, then filtered off again and dried *in vacuo* to give 1.57 g (20% yield) of the desired sulfone as a brown powder. This was used directly in the next reaction.

Step B


1.07 g (3.9 mmoles) of 5-phenylsulfonylindanone was taken up in dry THF (20 mL) at room temperature. A 1M solution of lithium aluminum hydride in THF (3.9 mL) was added slowly. The resulting mixture was stirred for one hour

- 23 -

- at room temperature, and quenched cautiously with 2M sodium hydroxide solution and then with water. Solid potassium carbonate was added, and the mixture was stirred until the salts formed a granular precipitate. This was filtered off and washed with ether, and the organic solvents were evaporated to give
- 5 0.998 g of the desired alcohol (92%), which was sufficiently pure for the next reaction.

Step C

- 0.700 g (2.55 mL) of the starting indanyl alcohol were taken up in neat
- 10 thionyl chloride (2 mL), and the resulting mixture was stirred at room temperature for one hour. The volatiles were removed on a rotary evaporator, and the crude chloride was taken up in DMF (5 mL) with N-hydroxyethylpiperazine (0.94 mL, 3 equiv.). This mixture was heated to about 60°C for
- 15 16 hours, cooled to room temperature, and poured into ice water (100 mL). The crude product was extracted with methylene chloride, the organic layers were dried and evaporated, and the residue was purified by column chromatography on silica gel. The column was eluted with ethyl acetate containing 5% methanol and then with ethyl acetate containing 5% methanol and 2% triethylamine to give the desired product.

20 Example 2:

- 24 -

6-Hydroxy-4-chromanone. 1:

A solution of resorcinol (11.0 g) in nitrobenzene (120 mL) at 5°C was treated dropwise with β -chloropropionyl chloride (12.7 g) with stirring.

Anhydrous AlCl_3 (31 g) was added portionwise with stirring, and the

- 5 temperature was kept below 15°C. After 15 minutes the mixture was heated to 40-45°C and kept there with stirring for 3 hours longer; it was then cooled to room temperature and allowed to stand overnight. The reaction mixture was poured with stirring into crushed ice (200 g) containing conc. HCl (10 mL). This was extracted with ether and the organic layer was extracted with 5% NaOH.
- 10 The basic layer was made acidic with conc. HCl. A tar-like solid separated out and was recrystallized from hot water to yield a white solid, m.p. 138-142°C.

7-(4'-Nitrophenoxy)-4-chromanone. 2:


A solution of 6-hydroxychromanone (1.64 g, 10 mmol) in DMF (10 mL) was treated with NaH (10 mmole, 60% in mineral oil) with stirring. After

15 30 minutes it was treated with a solution of 4-fluoronitrobenzene in DMF (5 ml) and stirred at 90°C for 6 hours. After cooling, it was diluted with ice water (75 mL) and then extracted with EtOAc (2 x 50 mL). Evaporation of the dried organic layer gave a gum-like residue (1.9 g). This crude product was used in the next step without further purification.

20 1-Cyclohexyl-4-[7-[(4-nitrophenyl)oxy]-chroman-4-yl]-piperazine. 3:

- Crude 7-(4'-nitrophenoxy)-4-chromanone (1.9 g) in EtOH (150 mL) was reduced with NaBH_4 (250 mg) at room temperature overnight. The crude product was purified through a column of silica gel ($\text{EtOAc}:\text{CH}_2\text{Cl}_2$, 2:8) to give pure 7-(4'-nitrophenoxy)-4-chromanol (550 mg). This compound (550 mg)
- 25 was converted to the chloride with thionyl chloride (0.28 g) in CH_2Cl_2 (30 mL) at 0°C for 1 hour and then at room temperature for 3 hours. Work-up yielded crude 4-chloro-7-(4'-nitrophenoxy)-chroman (500 mg). This was heated with N-cyclohexylpiperazine (0.9 g) at 130°C for one hour. The mixture was then diluted with water (35 mL), made basic with K_2CO_3 and extracted with EtOAc.
- 30 The residue from evaporation of the dried organic layer was purified through a column of TLC-grade silica gel (35 g). The desired product was recrystallized from acetonitrile, m.p. 67-68°C.

- 25 -

Example 3:**6-Hydroxytetralone, 4:**

- A solution of 6-methoxytetralone (35 mmol), and AlBr_3 (75 mmol) in toluene (250 mL) was heated on an oil bath at 100°C for 5 to 6 hours. The solution was cooled to room temperature and then poured onto a mixture of 1N HCl (200 mL) and crushed ice (500 g). This was stirred with EtOAc (300 mL) and filtered through a sintered glass funnel. The filtrate was transferred to a separatory funnel and the aqueous layer was extracted with EtOAc (2 x 100 mL). The organic layers were combined and dried with MgSO_4 , filtered and evaporated to a solid, 6-hydroxytetralone, which was pure by NMR and TLC (1:1 hexane:EtOAc).

6-[(2-Pyrimidinyl)oxy]-tetralone, 5:


- 6-Hydroxytetralone (31 mmol) was dissolved in dry DMF (50 mL), chilled in an ice/water bath and blanketed with a stream of nitrogen. NaH (60% in mineral oil, 31 mmol) was added slowly and in portions. Once gas evolution ceased, 2-chloropyrimidine (31 mmol) was added, the ice bath was removed and the solution heated at 100°C for 1.5 hours. It was then cooled to room temperature and the solvent was removed *in vacuo*. The residue was treated with water and CH_2Cl_2 (200 mL each). Evaporation of the organic layer yielded crude material which was purified by flash chromatography (1:1 hexane:EtOAc) to afford 6-[(2-pyrimidinyl)oxy]-tetralone as a light yellow powder.

4-Cyclohexyl-1-piperazinyl-6-[(2-pyrimidinyl)oxy]-1,2,3,4-tetrahydronaphthalene, 6:

- 6-[(2-Pyrimidinyl)oxy]-tetralone (3.2 mmol), N-cyclohexylpiperazine (3.2 mmol) and Ti(O-2-Pr)_4 (3.2 mmol) were dissolved in CH_2Cl_2 (10 mL). The solution was stirred at room temperature for 20 hours and then quenched with NaCNBH_3 (6.4 mmol in 5 mL EtOH). Water (20 mL) was added, the resulting

- 26 -

- mixture was filtered through a pad of 'Celite', and the residue was rinsed with CH_2Cl_2 (10 mL). The organic layer was dried with Na_2SO_4 and evaporated to yield the crude product as an oil. Purification by column chromatography on TLC-grade silica gel and 100:3:1 CH_2Cl_2 :EtOH: NH_4OH as eluant gave the pure product, 4-cyclohexyl-1-piperazinyl-[6-[(2-pyrimidinyl)oxy]]-1,2,3,4-tetrahydro-naphthalene. This was converted to the dimaleate salt, m.p. 179-181.5°C, by dissolving in EtOAc and heating with 2 equivalents of maleic acid.

Example 4:10 5-((2-Pyrimidinyl)oxy)indanone:

- 5-Hydroxyindanone (3.2 g, 21.6 mmol) was dissolved in dry DMF (20 mL) in a 3-neck flask fitted with an N_2 inlet and reflux condenser. The system was blanketed with N_2 , and 60% NaH in mineral oil (860 mg) was added slowly and in portions at 0°C. Then 2-chloropyrimidine (2.5 g) was added, the ice bath was replaced with an oil bath, and the solution was heated at 100°C for 3 hours. This was then cooled to room temperature before removal of DMF in vacuo. To the residue was added water (50 mL), the mixture was extracted with CH_2Cl_2 (50 mL), and the aqueous layer extracted twice more with CH_2Cl_2 . The organic layer was extracted with 1N NaOH (50 mL) to remove unreacted phenol and then dried with Na_2SO_4 and evaporated to a dark brown solid (3.6 g). The solid was extracted with EtOAc; filtration removed an insoluble precipitate. The filtrate was dried as before and evaporated to yield the product which was pure by NMR and TLC (1:1 hexane:EtOAc).

20 2-(2-(Cyclohexylmethyl)-3H-1-oxo-inden-5-yl)pyrimidine. 8:

- To piperidine (0.25 mL) at 0°C was added acetic acid (0.2 mL) followed by cyclohexanecarboxaldehyde (918 mg, 8.2 mmol) and 5-[(2-pyrimidinyl)oxy]-

- 27 -

indanone, 7 (1.85 g, 8.2 mmol) (synthesized by the same route as 5 above). The mixture was heated at 100°C for 25 minutes. Methanol (20 mL) was added to the hot solution, which, after cooling, was concentrated *in vacuo*. The residue was treated with water and CH₂Cl₂ (20 mL each), and the organic layer was dried with Na₂SO₄. Evaporation of the solvent gave the crude material as a dark brown oil. Purification by flash chromatography with 1:1 hexane:ethyl acetate resulted in the pure enone, 8.

2-(2-(Cyclohexylmethyl)-2,3-dihydro-1-oxo-indan-5-yl)pyrimidine:

The enone was cleanly reduced to the saturated ketone with 10% Pd-C catalyst (100 mg) on a Parr apparatus for 45 minutes. The catalyst was filtered off to yield the saturated ketone.

Alternative Reduction of Ketone (general procedure)

The ketone (1 equivalent) is dissolved in EtOH and NaBH₄ (0.75 equivalent) is added. This is stirred at room temperature and monitored by TLC (1:1 Hexane:EtOAc) until all ketone has disappeared (about 2 to 4 hours) and the reaction is complete. Then the EtOH is removed *in vacuo* and the residue treated with an equal amount of CH₂Cl₂ and water. The organic layer is dried with Na₂SO₄ and the solvent is evaporated off to yield the crude alcohol, which is used directly without purification.

Conversion of Alcohol to Chloride (general procedure)


The alcohol (1 equivalent) is dissolved in CH₂Cl₂ and chilled in an ice water bath. Thionyl chloride (1.2 equivalents) is then added and the solution stirred under a CaSO₄ drying tube and monitored by TLC (25% EtOAc in Hexane) until the alcohol has disappeared. An equivalent amount of water is added and the mixture is basified to pH 8 with solid NaHCO₃. The aqueous layer is extracted with CH₂Cl₂. The organic extracts are combined, dried and evaporated to provide the crude chloride, which is used without further purification.

2-[[1-(4-Cyclohexyl-1-piperaziny)-2-(cyclohexylmethyl)-2,3-dihydro-1H-inden-5-yl]oxy]pyrimidine, 9:

The chloride (1 equivalent) and cyclohexylpiperazine were dissolved in CH₃CN, refluxed for 2 hours, and then cooled to room temperature. Water was added and the mixture was extracted with EtOAc (4 x). The organic extracts were dried and evaporated to yield the crude product which was purified by column chromatography using TLC-grade silica gel and EtOAc as eluant.

- 28 -

2-[[1-(4-Cyclohexyl-1-piperazinyl)-2-(cyclohexylmethyl)-2,3-dihydro-1H-inden-5-yl]oxy]pyrimidine dimaleate was a mixture of diastereomers (proportions unknown), m.p. 150-154°C.

Example 5

5


The alcohol (0.4 g) was treated with thionyl chloride (20 mL) at ambient temperature for 3 hours. Thionyl chloride was removed under vacuum, the crude product was treated with piperidinopiperidine (1.22 g), and the reaction mixture heated overnight at 130°C. At the end of this time the reaction mixture was cooled to ambient temperature, diluted with dichloromethane and washed with 10% sodium hydroxide solution. The crude product was purified on silica gel (triethylamine:ethyl acetate 1:20) to give the desired product (0.05 g). HRMS for $C_{26}H_{35}N_2O_3S$: Calcd.: 455.2368; Found: 455.2373.

10

Example 6

2-[[1-(4-Cyclohexyl-1-piperazinyl)-2-(cyclohexylmethyl)-2,3-dihydro-1H-inden-5-yl]oxy]pyrimidine, 9:x

15



20

The chloride (0.6 g, 2.0 mmol) (from the first crude product of the process of Example 5), DMF (10 mL), sodium iodide (1.8 g), triethylamine (0.21 g) and piperazine (0.38 g, 2.0 mmol) were heated overnight at 50°C, and stirred for an additional 2 hours at 70°C. The reaction mixture was then cooled to room temperature and diluted with ethyl acetate. The solution was washed with 10% Na_2CO_3 , water and brine, and then concentrated to yield a mixture of two

- 29 -

diastereomers. Separation was carried out by chromatography on silica gel using ethyl acetate; diastereomer R_f: for Isomer A (compound 14, Table 1) = 0.3, and for Isomer B (compound 15, Table 1) = 0.4.


Further compounds that are described in the Tables or listed by formula
5 immediately after the Tables can be prepared by analogous methods.

While a number of embodiments of this invention are described herein, it is apparent that the embodiments can be altered to provide other embodiments that utilize the compositions and processes of this invention. Therefore, it will be appreciated that the scope of this invention includes alternative embodiments
10 and variations which are defined in the foregoing Specification; and the invention is not to be limited to the specific embodiments that have been presented herein by way of example.

- 30 -

Claims:

1. A compound having the structural formula



or a stereoisomer thereof, or a pharmaceutically acceptable salt, ester or solvate



- 5 thereof,
wherein:

Z is N, CH or C-alkyl;

X is -O-, -S-, -SO-, -SO₂-, -CO-, -CH₂-, -CONR²⁰-, -NR²⁰-SO₂-,
-NR²⁰CO-, or -SO₂-NR²⁰-;


- 10 Q is -O-, -S-, -SO-, -SO₂-, or -CH₂-;

R is



R¹ and R²¹ are independently selected from the group consisting of
hydrogen, alkyl, alkenyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl,
15 cycloalkenylalkyl, phenylalkyl, and hydroxyalkyl;

R² is cycloalkyl, cycloalkyl substituted with 1 to 3 independently selected
R³ groups, cycloalkenyl, cycloalkylalkyl,



(wherein R^Y is H, alkyl, alkenyl, SO₂R^Z or COR^Z
wherein R^Z is alkyl, alkenyl, aryl, heteroaryl, or cycloalkyl), with the proviso that

- 20 R² is R³-substituted-1-piperidinyl only when Z is CH or C-alkyl; or, when Z is

- 31 -

CH, R² may also be alkoxycarbonyl, -N(R⁹)(hydroxyalkyl) wherein R⁹ is H, hydroxyalkyl, or alkyl, or -N(R⁹)₂ wherein the two R⁹ groups may be joined to form an alkylene group;

5 R³, R⁴, R⁵, R⁶, R²², R²⁴, and R²⁵ are independently selected from the group consisting of H, alkyl, halo, alkoxy, benzyloxy, benzyloxy substituted by nitro or aminoalkyl, polyhaloalkyl, nitro, sulfonyl, hydroxy, amino, alkylamino, formyl, alkylthio, acyloxy, alkylsulfonyl, arylsulfonyl, acyl, alkoxycarbonyl, alkylsulfinyl, -OCONH₂, -OCONH-alkyl, -OCON(alkyl)₂, -NHCOO-alkyl, -NHCO-alkyl, phenyl, hydroxyalkyl, and 1-morpholinyl;

10 R⁸ is hydrogen, lower alkyl or cyclopropyl;



R²⁰ is H, phenyl or alkyl; and

R²⁷ and R²⁸ are independently selected from the group consisting of H, alkyl, hydroxyalkyl, alkoxyalkyl, arylalkyl, mercaptoalkyl, alkylthioalkyl, and carboxyalkyl, and additionally R²⁷ and R²⁸ may be joined to form an alkylene group; and

15 n is 0 or an integer from 1 to 3.

2. A compound of claim 1, wherein Z is N, Q is -CH₂- and n is 0, 1 or 2.

3. A compound of any of claims 1 or 2, wherein X is O, SO or SO₂, and


20 wherein R is  or , wherein R⁵ and R⁶ are H, CH₃, nitro, NH₂, acetylamino, or methoxy.

4. A compound of any of claims 1, 2 or 3, wherein R¹ is H, cycloalkyl, cycloalkylalkyl or alkyl and R²¹ is H.


25

5. A compound of any of claims 1, 2, 3 or 4, wherein X is O or SO₂, R² is cyclohexyl or 1-piperidinyl, and wherein at least one of R²⁷ and R²⁸ is alkyl and the other is H or alkyl.

6. A compound of claim 1 selected from the group consisting of the following
30 formulae:



- 32 -



wherein the methyl group attached to the piperazine ring is in the (R)-
5 conformation.

7. A pharmaceutical composition which comprises a compound as defined in claim 1 in combination with a pharmaceutically acceptable carrier.
8. A process for the preparation of a pharmaceutical composition as
10 defined in claim 7 comprising admixing a compound as claimed in any of claims 1 to 6 with a pharmaceutically acceptable carrier.
9. The use of a compound of any of claims 1 to 6 for the preparation of a medicament for treating a cognitive or neurodegenerative disease.
- 15 10. A kit for treating a cognitive or neurodegenerative disease comprising separate containers in a single package that contain pharmaceutical compounds for use in combination, wherein one container contains a compound in accordance with any of claims 1 to 6 that enhances acetylcholine release and a separate container contains an acetylcholinesterase inhibitor, said compound

– 33 –

and inhibitor each being in a pharmaceutically acceptable carrier and their combined quantities being an effective amount.

11. A method for treating a cognitive or neurodegenerative disease comprising administering to a patient suffering from said disease an effective amount of a compound of any of claims 1 to 6, alone or in combination with an acetylcholinesterase inhibitor.
- 5

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US 97/10696

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D295/096 C07D311/68 C07D213/643 C07D405/12 A61K31/495
A61K31/44 A61K31/35

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	C. MELCHIORRE ET AL.: JOURNAL OF MEDICINAL CHEMISTRY, vol. 36, no. 23, 1993, pages 3734-7, XP002048425 cited in the application see page 3734, left-hand column, compound 3 ---	1
A	WO 94 12493 A (SANDOZ) 9 June 1994 see claims 2,3,9 ---	1,7
A	EP 0 028 449 A (BEECHAM GROUP LTD.) 13 May 1981 see claims 1,4,11 ---	1,7
A	EP 0 321 175 A (BEECHAM GROUP PLC) 21 June 1989 see claims 1,14; examples ---	1,7
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "B" document member of the same patent family

Date of the actual completion of the international search

27 November 1997

Date of mailing of the international search report

09.12.97

Name and mailing address of the ISA

European Patent Office, P.O. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hass, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/10696

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 92 10192 A (H. LUNDBECK A/S) 25 June 1992 see claims 1,6 ---	1,7
A	DE 24 12 522 A (F. HOFFMANN-LA ROCHE & CO. AG) 10 October 1974 see claims 13,14 ---	1,7
A	WO 93 08799 A (SMITHKLINE BEECHAM CORP.) 13 May 1993 cited in the application see claims 1,6 ---	1,7
A	J. M. EVANS ET AL.: JOURNAL OF MEDICINAL CHEMISTRY, vol. 27, no. 9, 1984, pages 1127-31, XP002048426 see page 1129, table II ---	1
A	W. LOGEMANN ET AL.: BRITISH JOURNAL OF PHARMACOLOGY, vol. 17, 1961, pages 286-96, XP000576553 cited in the application -----	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/10696

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9412493 A	09-06-94	CA 2122494 A AU 665040 B AU 3082692 A EP 0623129 A FI 943489 A JP 7503487 T NO 942744 A PL 172371 B SK 88994 A	09-06-94 14-12-95 22-06-94 09-11-94 22-09-94 13-04-95 22-07-94 30-09-97 07-06-95
EP 28449 A	13-05-81	JP 56057786 A US 4366163 A	20-05-81 28-12-82
EP 321175 A	21-06-89	AU 2678088 A DK 691488 A JP 2000141 A	22-06-89 15-06-89 05-01-90
WO 9210192 A	25-06-92	AT 121294 T AU 653375 B AU 8959991 A CA 2097715 A DE 69109132 D DE 69109132 T EP 0560868 A ES 2071479 T HK 122995 A IL 100164 A JP 6503073 T NO 179947 B NZ 240734 A US 5643784 A	15-05-95 29-09-94 08-07-92 05-06-92 24-05-95 31-08-95 22-09-93 16-06-95 04-08-95 18-06-96 07-04-94 07-10-96 25-02-94 01-07-97
DE 2412522 A	10-10-74	CH 593970 A AR 218600 A AT 344177 B AT 344181 B AT 344182 B AU 6582774 A CA 1024516 A DD 113545 A	30-12-77 30-06-80 10-07-78 10-07-78 10-07-78 21-08-75 17-01-78 12-06-75

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 97/10696

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 97/10696

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

Claims Nos.: 11

because they relate to subject matter not required to be searched by this Authority, namely:

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

Remark : Although claim 11 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/10696

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 2412522 A		FR 2223027 A	25-10-74
		GB 1464977 A	16-02-77
		GB 1464978 A	16-02-77
		GB 1464979 A	16-02-77
		JP 49135982 A	27-12-74
		LU 69734 A	04-02-76
		NL 7404312 A	02-10-74
		NL 7606088 A	08-12-76
		US 3954764 A	04-05-76
		US 4044010 A	23-08-77
		BE 813003 A	30-09-74
		CH 593971 A	30-12-77
		CH 593969 A	30-12-77
		SE 403290 B	07-08-78
		ZA 7400940 A	26-02-75
<hr/>			
WO 9308799 A	13-05-93	AP 433 A	17-11-95
		AU 669866 B	27-06-96
		AU 3125993 A	07-06-93
		BG 98752 A	31-05-95
		BR 9206722 A	18-07-95
		CN 1073161 A	16-06-93
		CN 1145223 A	19-03-97
		CZ 9401109 A	18-01-95
		EP 0612244 A	31-08-94
		FI 942059 A	04-07-94
		HU 67665 A	28-04-95
		JP 7501322 T	09-02-95
		MX 9206348 A	30-06-94
		NO 941650 A	01-07-94
		NZ 245000 A	28-05-96
		OA 9921 A	15-08-94
		PT 101038 A	28-02-94
		SK 52194 A	08-03-95
		ZA 9208467 A	05-05-93
<hr/>			